

Single Transverse-Spin Asymmetry in $p^\uparrow p \rightarrow \pi X$ and $ep^\uparrow \rightarrow \pi X$

Y. Koike

Dept. of Physics, Niigata University, Japan

Our interest in the single transverse spin asymmetry for the pion production in pp and ep collisions resides in the fact that it probes particular quark-gluon correlation in the hadrons (higher twist effect) which is not included in the twist-2 parton distributions. QCD analysis for the asymmetry is crucial for the ongoing RHIC-SPIN, HERMES and COMPASS experiments.

In this contribution, we will present an analysis for the asymmetry in the framework of the collinear factorization [1-2]. According to the generalized QCD factorization theorem, the polarized cross section for $p^\uparrow p \rightarrow \pi X$ consists of three kinds of twist-3 cross sections,

$$(A) \quad G_a(x_1, x_2) \otimes q_b(x') \otimes \hat{q}_{c \rightarrow \pi}(z) \otimes \hat{\sigma}_{ab \rightarrow c}^1 \quad (1)$$

$$(B) \quad \delta q_a(x) \otimes E_b(x_1, x_2) \otimes \hat{q}_{c \rightarrow \pi}(z) \otimes \hat{\sigma}_{ab \rightarrow c}^2 \quad (2)$$

$$(C) \quad \delta q_a(x) \otimes q_b(x') \otimes \hat{E}_{c \rightarrow \pi}(z_1, z_2) \otimes \hat{\sigma}_{ab \rightarrow c}^3 \quad (3)$$

and the one for $ep^\uparrow \rightarrow \pi X$ likewise receives two contributions,

$$(A') \quad G_a(x_1, x_2) \otimes \hat{q}_{a \rightarrow \pi}(z) \otimes \hat{\sigma}, \quad (4)$$

$$(C'') \quad \delta q_a(x) \otimes \hat{E}_{a \rightarrow \pi}(z_1, z_2) \otimes \hat{\sigma}'. \quad (5)$$

Here the functions with two variables (momentum fractions) $G_a(x_1, x_2)$, $E_a(x_1, x_2)$, $\hat{E}_{c \rightarrow \pi}(z_1, z_2)$ are twist-3 quantities: G_a and E_a are respectively, the transversely polarized distribution and the unpolarized distribution functions in the nucleon. $\hat{E}_{c \rightarrow \pi}$ is the unpolarized fragmentation function for the pion. a , b and c stand for the parton's species. Other functions are twist-2; $q_b(x)$ the unpolarized distribution, $\delta q_a(x)$ the transversity distribution, $\hat{q}_{c \rightarrow \pi}$ the unpolarized fragmentation function. $\hat{\sigma}_{ab \rightarrow c}^1$ etc. represents the partonic cross section which yields large transverse momentum of the pion.

The analyses of (A) and (B) contributions have been provided in [1] and [2], respectively. Here we extend the analysis to the (C) term (for a preliminary study, see [3]) and the ep^\uparrow case so that we can have more thorough understanding on the hadron structure and the reaction mechanism.

References

- [1] J. Qin and G. Sterman, Phys. Rev. **D59** (1999) 014004.
- [2] Y. Kanazawa and Y. Koike, Phys. Lett. **B478** (2000) 121; **B490** (2000) 99.
- [3] Y. Koike, hep-ph/0106260, Proceedings of DIS2001. (Bologna, Italy, April 2001.)